



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Then $AM + CB = AM + PM = l$. Likewise, $AM' + C'B' = l$. There will be two solutions, one solution, or no solution according as PQ cuts the circle, is tangent to the circle, or lies outside of the circle. In other words the problem is possible only when $2r < l < (1 + \sqrt{5})r$.

Also solved analytically, by H. C. FEEMSTER.

413. Proposed by DAVID R. KELLEY, New York City.

To construct a triangle, given the base, the vertical angle, and the ratio of the altitude from the vertex on the base to the difference of the other two sides.

SOLUTION BY THE PROPOSER.

Let ABC be the required triangle. Then since side AB and $\angle C$ are known, we can construct the circumcircle of the triangle. Let the bisectors of the interior angle C and the exterior angle C meet the circle in D and E , respectively. Then D and E are fixed points, and ED is the diameter of the circle, and AE is fixed. Draw EG perpendicular to BC produced. Then $CG = \frac{1}{2}(AC - BC)$. Hence, if CP be the altitude from C on AB , CG/CP is also known, since by hypothesis $(AC - CB)/CP$ is the given ratio. Again, since in the right triangle ECG , $\angle GCE$ is known, therefore GC/EC is known, and therefore, by above, EC/CP is known. Now let EC meet AB in F . Then triangles DEC and CFP are similar. Hence, $EC/CP = DE/CF$. But EC/CP is given. Hence DE/CF is also known, and therefore, since DE is the diameter of the circumcircle of triangle ABC , CF can be found. Also $EC \cdot EF = EA^2$. Hence, since CF and EA are known, EC can be constructed and hence point C can be found. See Fig. 2.

Also solved by H. C. FEEMSTER, C. N. SCHMALL, and A. M. HARDING.

414. Proposed by H. C. FEEMSTER, York, Neb.

To construct the cyclic quadrilateral, having given its four sides.

SOLUTION BY THE PROPOSER.

Construction.—Let the lengths of the given sides be a, b, c, d . Draw $AD = d$, and produce it to E so that $DE : c = a : b$. Now describe a circle which is the locus of a point P such that $EP : PA = c : b$ (circle of Apollonius). Describe also a circle with the center D and radius c . Let C be one point of intersection of these circles. Then on AC as a base construct a triangle ABC similar to EDC . Then $ABCD$ is the quadrilateral required. See Fig. 3.

Proof.—By construction $AD = d$, $DC = c$. Now since the angles ABC and EDC are equal, therefore $\angle ABC$ and $\angle ADC$ are supplementary and the figure $ABCD$ is cyclic. It now remains to prove that $AB = a$, and $BC = b$.

By construction, $ED : c = a : b$, or $ED : DC = a : b$. Also by construction $EC : CA = c : b$, or $EC : DC = CA : b$. Hence

$$ED : DC : EC = a : b : CA. \quad (1)$$

But by similar triangles, we have

$$ED : DC : EC = AB : BC : CA. \quad (2)$$

Then by (1) and (2), $AB = a$, $BC = b$.